

Electron Tagging

W Emulsion, SFT, EMCal

Bruce Baller


1/30/03

Goal & Method

- Use information from all available systems to tag electrons
- Electron tagging in emulsion is 90% (BB code) efficient for > 1 radiation length and momentum > 10 Gev
 - Efficiency for electrons in period 4 events is only 50%
- Developed SFT and EMCal electron tag code for located events
- Emulsion electron tag code is unchanged since last report
- Developed voting scheme which combines Emulsion (EML), SFT (SFT) and EMCal (EMC) tagging results to improve efficiency and reduce false positives

Radiation length cuts

- EMC and SFT taggers use cuts based on accumulated radiation length in each station
- Use variant of Byron's code in new routine `getradlen.sf`



- Returns array of radiation length in each SFT station & EMCal
 - `Radstn(4), emcrad`
- Ex: `radstn = (0, 1, 3.1, 5.1)` `emcrad= 6.1`

EMC tagger


- Define a window ($\text{win} = 20 \text{ cm}$) on the face of the calorimeter for matching primary tracks to clusters
- Only consider primary tracks separated by $\delta\theta > \text{win}/(z_{\text{cal}} - z_{\text{vtx}})$ from other tracks
- Require tracks project to face of calorimeter and $n_{\text{seg}} > 2$
- Set a track flag bit to indicate that track passes EMC tagger cuts
- Define min cluster energy cut
 - $E_{\text{cut}} = 0.8 \cdot (7 - \text{emcrad})$
 - $\text{Min } e_{\text{cut}} = 0.3$
- Find closest cluster to track projection with $e_{\text{clus}} > e_{\text{cut}}$ and $\delta r < 20 \text{ cm}$
- Set “EMC_TAG” track flag bit
- Store cluster energy in track array

SFT Tagger

- Consider tracks with $nseg > 2$
- Only consider primary tracks separated by $\delta\theta > 40$ mr from other primary tracks
- Set a track flag bit to indicate that track passes SFT tagger track cuts
- Define a window for each SFT station for summing pulse height
 - EM shower width \sim radiation length
 - Determine scale factor by scanning MC events to maximize pulse height and minimize pulse height cross-talk between tracks
 - $Cut = 0.002 * radstn(stn)$
- Sum PH in each view (X,U,V) within cut
- Count number of planes traversed
- Normalize PH to MIP's/plane traversed
- Correct PH in X plane (4 fiber planes)

SFT window cut for 29 GeV electron

$\langle PH \rangle = 20$ MIP's/plane



SFT Tagger - Cont

- Determine PH difference in the views
 - Eliminates false positives - hadron tracks with overlapping EM shower in one view
- $\langle PH \rangle = (PHX + PHU + PHV)/3$
- $PH\ rms = [(PHX - PHU)^2 + (PHX - PHV)^2 + (PHU - PHV)^2]^{1/2} / \langle PH \rangle$
- “SFT_TAG” = $\langle PH \rangle > 10\text{ MIP's/plane}$ & $PH\ rms < 0.5$
- Set “SFT_TAG” track flag bit
- Store $\langle PH \rangle$ and $PH\ rms$ in track array
- One ****could**** correct PH from overlapping showers in one view using information in the other views

Combining Tagger Results

- EML/SFT are complementary to EMCal
 - Brehm tracks in emulsion \rightarrow large PH in SFT
 \rightarrow reduced EMCal energy
 - Energy sharing dependent on emulsion target station & radiation length
- Define ALL tagger which uses EML, SFT and EMC tagger results
- Ordered set of cuts
- $ALL = \overline{EML_TAG}$
- $ALL = \overline{ALL} \bullet SFT_TAG \bullet EMC_TAG$
- $ALL = \overline{ALL} \bullet EMC_TAG \bullet STN > 2$
- $ALL = \overline{ALL} \bullet (STN > 1 \bullet STN < 4) \bullet$
 $\langle PH \rangle > 15 \text{ MIP/PLANE}$

MC Tagger Results

- Evaluate efficiency and rate of false positives on 500 electron CC events in Period 4
- Define track class
 - 0 = True electron failed all tagger track selection cuts
 - 1 = True electron correctly tagged
 - 2 = True electron tagged as hadron
 - 3 = True hadron tagged as electron
 - 4 = True hadron tagged as hadron
 - 5 = True hadron failed all tagger track selection cuts

MC Tagger Results

Track class

	True Electrons 472			True Hadrons 1330		
Tagger	0	1	2	3	4	5
ALL	14%	72%	14%	5%	58%	37%
EMC	28%	51%	21%	6%	45%	48%
SFT	16%	62%	21%	8%	69%	24%
EML	1%	51%	47%	1%	98%	2%

- 72% of electrons correctly tagged
- 5% of hadrons tagged as electrons
- Other bits of information
- 12% of events have >1 electron attached to the primary ($IP < 5$ micron)
- The true primary electron is tagged in 86% of electron CC events

What next?

- Check failed true electron tags
 - I checked 23 class 2 events with $\text{evt_wght} > 30$
 - 18 event tracks have true electron energy < 10 GeV
 - 21 tracks have no brehm tracks visible in the emulsion
 - 12 tracks failed the SFT tagger 10 MIP cut
 - 9 tracks missed the EMCal
 - 4 tracks have no EMCal cluster < 20 cm
 - 9 tracks failed the EMCal ecut ($E_{\text{clus}} < 1$ GeV)
- Possible improvements
 - Not much room for improvement in EMC or EML
 - SFT: Unfold overlapping showers
- Send tagging results to Nonaka?