

## $\chi_{b0}(2P)$

$I^G(JPC) = 0^+(0^{++})$   
 $J$  needs confirmation.

Observed in radiative decay of the  $\Upsilon(3S)$ , therefore  $C = +$ . Branching ratio requires E1 transition, M1 is strongly disfavored, therefore  $P = +$ .

### $\chi_{b0}(2P)$ MASS

VALUE (GeV)	DOCUMENT ID	TECN	COMMENT
<b><math>10.2321 \pm 0.0006</math> OUR AVERAGE</b>			
$10.2312 \pm 0.0008 \pm 0.0012$	<sup>1</sup> HEINTZ 92	CSB2	$e^+ e^- \rightarrow \gamma X, \ell^+ \ell^- \gamma\gamma$
$10.2323 \pm 0.0007$	<sup>2</sup> MORRISON 91	CLE2	$e^+ e^- \rightarrow \gamma X$

<sup>1</sup> From the average photon energy for inclusive and exclusive events and assuming  $\Upsilon(3S)$  mass =  $10355.3 \pm 0.5$  MeV. Supersedes HEINTZ 91 and NARAIN 91.  
<sup>2</sup> From  $\gamma$  energy below assuming  $\Upsilon(3S)$  mass =  $10355.3 \pm 0.5$  MeV. The error on the  $\Upsilon(3S)$  mass is not included in the individual measurements. It is included in the final average.

### $\gamma$ ENERGY IN $\Upsilon(3S)$ DECAY

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>122.8 \pm 0.5</math> OUR AVERAGE</b> Error includes scale factor of 1.1.				
123.0 $\pm$ 0.8	4959	<sup>3</sup> HEINTZ 92	CSB2	$e^+ e^- \rightarrow \gamma X$
124.6 $\pm$ 1.4	17	<sup>4</sup> HEINTZ 92	CSB2	$e^+ e^- \rightarrow \ell^+ \ell^- \gamma\gamma$
$122.3 \pm 0.3 \pm 0.6$	9903	MORRISON 91	CLE2	$e^+ e^- \rightarrow \gamma X$

<sup>3</sup> A systematic uncertainty on the energy scale of 0.9% not included. Supersedes NARAIN 91.  
<sup>4</sup> A systematic uncertainty on the energy scale of 0.9% not included. Supersedes HEINTZ 91.

### $\chi_{b0}(2P)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1 \quad \gamma \Upsilon(2S)$	( $4.6 \pm 2.1$ ) %
$\Gamma_2 \quad \gamma \Upsilon(1S)$	( $9 \pm 6$ ) $\times 10^{-3}$

### $\chi_{b0}(2P)$ BRANCHING RATIOS

$\Gamma(\gamma \Upsilon(2S))/\Gamma_{\text{total}}$	$\Gamma_1/\Gamma$
<b><math>&lt;0.089</math></b>	
<b><math>0.046 \pm 0.020 \pm 0.007</math></b>	$90$

<sup>5</sup> Using  $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.37 \pm 0.26)\%$ ,  $B(\Upsilon(3S) \rightarrow \gamma\gamma \Upsilon(2S)) \times 2 B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) < 1.19 \times 10^{-4}$ , and  $B(\Upsilon(3S) \rightarrow \chi_{b0}(2P)\gamma) = 0.049$ .  
<sup>6</sup> Using  $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.44 \pm 0.10)\%$ ,  $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) = (6.0 \pm 0.4 \pm 0.6)\%$  and assuming  $e\mu$  universality. Supersedes HEINTZ 91.

$\Gamma(\gamma \Upsilon(1S))/\Gamma_{\text{total}}$		$\Gamma_2/\Gamma$		
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.025	90	<sup>7</sup> CRAWFORD	92B CLE2	$e^+ e^- \rightarrow \ell^+ \ell^- \gamma\gamma$
<b>0.009±0.006±0.001</b>		<sup>8</sup> HEINTZ	92 CSB2	$e^+ e^- \rightarrow \ell^+ \ell^- \gamma\gamma$

<sup>7</sup> Using  $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.57 \pm 0.07)\%$ ,  $B(\Upsilon(3S) \rightarrow \gamma\gamma \Upsilon(1S)) \times 2 B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) < 0.63 \times 10^{-4}$ , and  $B(\Upsilon(3S) \rightarrow \chi_b0(2P)\gamma) = 0.049$ .

<sup>8</sup> Using  $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.57 \pm 0.07)\%$ ,  $B(\Upsilon(3S) \rightarrow \gamma\chi_b0(2P)) = (6.0 \pm 0.4 \pm 0.6)\%$  and assuming  $e\mu$  universality. Supersedes HEINTZ 91.

## $\chi_b0(2P)$ REFERENCES

CRAWFORD	92B	PL B294 139	G. Crawford, R. Fulton	(CLEO Collab.)
HEINTZ	92	PR D46 1928	U. Heintz <i>et al.</i>	(CUSB II Collab.)
HEINTZ	91	PRL 66 1563	U. Heintz <i>et al.</i>	(CUSB Collab.)
MORRISON	91	PRL 67 1696	R.J. Morrison <i>et al.</i>	(CLEO Collab.)
NARAIN	91	PRL 66 3113	M. Narain <i>et al.</i>	(CUSB Collab.)

## OTHER RELATED PAPERS

EIGEN	82	PRL 49 1616	G. Eigen <i>et al.</i>	(CUSB Collab.)
HAN	82	PRL 49 1612	K. Han <i>et al.</i>	(CUSB Collab.)