

## § 529.2464

(3) Milk that has been drawn from animals within 48 hours of such treatment may not be used for food.

[41 FR 10984, Mar. 15, 1976, as amended at 43 FR 29290, July 7, 1978; 55 FR 29842, July 23, 1990; 55 FR 31481, Aug. 2, 1990; 62 FR 8372, Feb. 25, 1997]

### § 529.2464 Ticarcillin powder.

(a) *Specifications.* Each vial contains ticarcillin disodium equivalent to 6 grams of ticarcillin to be reconstituted with 25 milliliters of sterile water for injection or sterile physiological saline.

(b) *Sponsor.* See No. 000069 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount.* 6 grams per day, intrauterine, for 3 consecutive days during estrus.

(2) *Indications for use. Horses.* Intrauterine treatment of endometritis caused by beta-hemolytic streptococci.

(3) *Limitations.* For intrauterine use in horses only. Infuse aseptically. Not for use in horses raised for food production. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37336, Aug. 18, 1992, as amended at 60 FR 55660, Nov. 2, 1995]

### § 529.2503 Tricaine methanesulfonate.

(a) *Chemical name.* Ethyl-*m*-amino-benzoate methanesulfonate.

(b) *Sponsor.* See Nos. 050378 and 051212 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) It is used for the temporary immobilization of fish, amphibians, and other aquatic cold-blooded animals (poikilotherms) as an aid in handling during manual spawning (fish stripping), weighing, measuring, marking, surgical operations, transport, photography, and research.

(2) It is used as follows:

(i) For fish the drug is added to ambient water at a concentration of from 15 to 330 milligrams per liter depending upon the degree of anesthetization or sedation desired, the species and size of the fish, and the temperature and softness of the water. Preliminary tests of solutions must be made with small numbers of fish to determine the desired rates of sedation or anesthesia and the appropriate exposure times for

## 21 CFR Ch. I (4-1-01 Edition)

the specific lots of fish under prevailing conditions.

(ii) For amphibians and other aquatic coldblooded animals, the drug is added to ambient water in concentrations of from 1:1000 to 1:20,000 depending upon species and stage of development.

(iii) Do not use within 21 days of harvesting fish for food. Use in fish intended for food should be restricted to Ictaluridae, Salmonidae, Esocidae, and Percidae, and water temperature should exceed 10° C. (50° F.). In other fish and in cold-blooded animals, the drug should be limited to hatchery or laboratory use.

[40 FR 13881, Mar. 27, 1975, as amended at 49 FR 5748, Feb. 15, 1984; 51 FR 11439, Apr. 3, 1986; 63 FR 7702, Feb. 17, 1998]

## PART 530—EXTRALABEL DRUG USE IN ANIMALS

### Subpart A—General Provisions

Sec.

- 530.1 Scope.
- 530.2 Purpose.
- 530.3 Definitions.
- 530.4 Advertising and promotion.
- 530.5 Veterinary records.

### Subpart B—Rules and Provisions for Extralabel Uses of Drugs in Animals

- 530.10 Provision permitting extralabel use of animal drugs.
- 530.11 Limitations.
- 530.12 Labeling.
- 530.13 Extralabel use from compounding of approved new animal and approved human drugs.

### Subpart C—Specific Provisions Relating to Extralabel Use of Animal and Human Drugs in Food-Producing Animals

- 530.20 Conditions for permitted extralabel animal and human drug use in food-producing animals.
- 530.21 Prohibitions for food-producing animals.
- 530.22 Safe levels and analytical methods for food-producing animals.
- 530.23 Procedure for setting and announcing safe levels.
- 530.24 Procedure for announcing analytical methods for drug residue quantification.
- 530.25 Orders prohibiting extralabel uses for drugs in food-producing animals.