

Jet quenching on hadron yields and ratios at RHIC and LHC

P. Levai^a G.G. Barnafoldi^a G. Papp^b G. Fai^c Y. Zhang^c
M. Gyulassy^d

^a*RMKI Research Institute for Particle and Nuclear Physics, Budapest, Hungary*

^b*HAS Research Group for Theoretical Physics, Eötvös University, Budapest,
Hungary*

^c*CNR, Department of Physics, Kent State University, Kent, OH, USA*

^d*Physics Department, Columbia University, New York, NY, USA*

Presented by: P. Levai

Abstract

The production and interaction of jets dominates physical observables in the high transverse-momentum region in ultrarelativistic heavy-ion collisions. One of these observables is the energy loss of high-energy jets ("jet quenching"). This phenomenon is an excellent diagnostic tool to investigate the properties of hot and dense matter. In a new calculation the energy loss was determined in the realistic case of a few scattering centers ($N_{scat} = 1, 2, 3, \dots$), and found a quadratic dependence on the opacity (L/λ) of the surrounding matter. The results of this calculation are applied in a perturbative QCD based study of pion and kaon production in heavy-ion collisions at RHIC ($\sqrt{s} = 56, 130$ A GeV). Absolute hadron yields display a close to linear shift with increasing energy loss. However, certain particle ratios (K^+/π^+ , K^-/K^+) show detectable modifications due to jet quenching. We also discuss our results at LHC energy.
